

**Bainbridge, Georgia:  
Later Maturity Cotton Variety Performance, 2008, Irrigated**

Variety	Lint Yield lb/acre	Lint %	Uniformity		Strength* g/tex	Micronaire* units
			Index*	Length* inches		
PHY375WRF	<b>1762</b>	44.3	83.0	1.18	32.3	3.8
BCSX0727B2RF	<b>1727</b>	43.8	83.0	1.18	33.9	4.4
DP 0935 B2RF	<b>1655</b>	44.4	82.6	1.15	33.3	4.3
DP174RF	1589	46.6	84.1	1.22	32.9	4.3
PHY485WRF	1560	42.4	83.3	1.15	36.1	4.3
BCSX0102LLB2	1536	41.3	84.8	1.25	36.3	3.8
GA2004358	1523	43.3	83.4	1.20	35.5	4.2
DP 555 BG/RR	1521	46.2	81.8	1.15	33.7	4.4
GA2004371	1511	45.9	83.5	1.20	34.1	4.7
DP161B2RF	1508	41.8	83.7	1.23	35.6	4.2
PHY480WR	1492	40.4	85.3	1.22	36.0	4.6
BCSX0721B2RF	1472	43.9	84.1	1.20	32.9	4.2
PHY440W	1451	41.1	83.1	1.18	33.7	3.8
PHY425RF	1444	42.6	84.0	1.18	37.1	4.8
GA2004137	1428	44.6	83.2	1.18	35.5	4.9
GA2004392	1407	41.8	83.0	1.18	36.0	5.0
BCSX0614B2RF	1306	40.3	82.7	1.20	33.6	3.6
DP 164 B2RF	1264	41.9	83.3	1.21	35.8	4.1
Average	1509	43.1	83.4	1.19	34.7	4.3
LSD 0.10	166	1.2	1.2	0.03	1.8	0.5
CV %	9.3	2.5	0.8	1.41	2.9	6.6

\* A random quality sample was taken on the picker during cotton harvest.

**Bolding** indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Planted: May 14, 2008.

Harvested: October 7, 2008.

Data as of 12/03/08.