

**Midville, Georgia:
Later Maturity Cotton Variety Performance, 2008, Irrigated**

Variety	Lint Yield lb/acre	Lint %	Uniformity		Strength* g/tex	Micronaire* units
			Index*	Length* inches		
PHY480WR	2291	42.2	84.7	1.19	29.2	4.4
DP 555 BG/RR	2255	45.0	82.0	1.17	30.6	4.1
DP 0935 B2RF	2228	44.1	83.9	1.16	27.4	4.1
DP161B2RF	2187	43.3	84.5	1.24	32.2	3.7
PHY440W	2161	42.8	83.8	1.17	29.8	4.1
BCSX0727B2RF	2136	43.9	83.6	1.17	27.7	4.4
DP174RF	2120	46.2	84.5	1.24	27.1	4.2
PHY375WRF	2010	42.7	83.2	1.20	27.3	4.0
BCSX0102LLB2	2006	40.7	85.2	1.27	32.3	3.7
PHY485WRF	1987	41.7	84.3	1.19	29.5	4.4
GA2004358	1934	42.4	83.1	1.20	31.6	4.5
GA2004392	1897	41.7	85.3	1.18	29.7	4.8
BCSX0614B2RF	1872	39.5	84.9	1.24	29.6	4.2
PHY425RF	1794	40.7	85.1	1.20	29.2	4.3
GA2004137	1740	43.4	84.4	1.20	30.7	4.4
DP 164 B2RF	1706	42.1	83.6	1.21	31.6	4.1
BCSX0721B2RF	1621	46.9	83.4	1.21	27.9	4.2
GA2004371	1570	44.7	85.6	1.19	29.6	4.7
Average	1973	43.0	84.1	1.20	29.6	4.2
LSD 0.10	230	0.9	1.8	0.03	1.8	0.3
CV %	9.8	1.8	1.0	0.28	3.6	3.7

* A random quality sample was taken on the picker during cotton harvest.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Planted: First planting on May 1, 2008; due to strong wind and blowing sand damage after emergence, replanted on May 28, 2008.

Harvested: October 22, 2008.

Data as of 11/19/08.